

# PAPILIO AMPHIMEDON.

VAR.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

## CHARACTER SPECIFICUS, &c.

PAPILIO alis concoloribus fuscis: anterioribus albo radiatis; posterioribus macula quinque-fida rubra lunulisque albis.

*Fabr. spec. ins.* 2. p. 8.

EQ. TR.

*Var?*

Alis superioribus fuscis ochraceo radiatis, inferioribus flavis nigro maculatis.

*Seb. mus.* 4. p. 21. t. 16. f. 7. 8.

---

Inter maximos papiliones exoticos numeratur Papilio Amphimedon. In insula Amboina generatur, & magnitudine vera in tabula exprimitur.

# AMPHIMEDON.

VAR.



## GENERIC CHARACTER.

*Antennæ* commonly thickening towards the end into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight diurnal.)

## SPECIFIC CHARACTER, &c.

Brown Butterfly, with the upper wings radiated with white; the lower with a five-cleft red spot and white marginal crescents.

*Var* ?

Upper wings radiated with ochre-color; lower yellow with black variegations.

---

This insect is one of the largest of the exotic Butterflies, and is a native of Amboina. The plate represents it in its natural size.

*London. Published & sold by W. Woodcock, Strand, near St. Dunstons Church.*

